

## Daily Soda Or SSB Consumption (School-age)

### Daily Soda Or SSB Consumption (School-age)

- Most recent data (2014): 17% of King County youth attending public schools in the 6th,8th,10th and 12th grades.
- Trend over time: 2004-14: falling
- Description: Drank non-diet soda or other sugar-sweetened beverage (SSB) once or more daily
- Healthy People 2020 Goal: None listed for this indicator.

Daily soda or SSB consumption (school-age), King County, 2012 and 2014 average

|   | Percent | Lower CI | Upper CI |
|---|---------|----------|----------|
| <b>King County</b> (school-age in 6th, 8th, 10th and 12th grades) | 23      | 22       | 24       |
| <b>GRADE</b>  |         |          |          |
| Grade 6   | 24      | 23       | 26       |
| Grade 8   | 23      | 21       | 24       |
| Grade 10  | 23      | 22       | 25       |
| Grade 12  | 23      | 21       | 24       |
| <b>GENDER</b>   |         |          |          |
| Male  | 27      | 27       | 28       |
| Female  | 19      | 18       | 20       |
| <b>RACE/ETHNICITY</b>   |         |          |          |
| AIAN  | 33      | 30       | 37       |
| Asian   | 18      | 17       | 20       |
| Black   | 29      | 28       | 31       |
| Hispanic  | 31      | 30       | 33       |
| Multiple  | 23      | 22       | 25       |
| NHPI  | 30      | 27       | 33       |
| White   | 22      | 21       | 22       |
| Other   | 24      | 23       | 26       |
| <b>REGION</b>   |         |          |          |
| East  | 20      | 19       | 21       |
| North   | 20      | 19       | 22       |
| Seattle   | 21      | 20       | 24       |
| South   | 27      | 26       | 28       |
| <b>STATE</b>  |         |          |          |
| Washington State  | 26      | 25       | 27       |

#### Comparisons:

Higher than King County rate (yellow)

Lower than King County rate (green)

#### Notes:

Source: Healthy Youth Survey.

Prepared by Public Health - Seattle & King County, APDE, 08/2016.


CI is 95% Confidence Interval.

\* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.

Data by income or poverty level not available.

## Daily soda or SSB consumption (school-age) King County, 2012 and 2014 average



Source: Healthy Youth Survey.

Prepared by Public Health - Seattle & King County, APDE, 08/2016.


[-----] Confidence interval shows range that includes true value 95% of the time.

\* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.


Data by income or poverty level not available.

**Daily soda or SSB consumption (school-age) by regions  
King County, 2012 and 2014 average**



Source: Healthy Youth Survey.  
Prepared by Public Health - Seattle & King County, APDE, 08/2016.  
N/A: Not available.

# Daily soda or SSB consumption (school-age) King County, 2004-2014



Trends over time: King County (2004-14: falling); East KC (2004-14: falling); North KC (2004-14: falling); Seattle (2004-14: falling);  
 South KC (2004-14: falling); Washington State (2004-14: falling)  
 Prepared by Public Health - Seattle & King County, APDE, 08/2016.  
 Source: Healthy Youth Survey.  
 South KC (2004-14: falling); Washington State (2004-14: falling)

Produced by the  
[Assessment, Policy Development & Evaluation Unit](#)  
[Public Health - Seattle & King County](#)  
 08/16